

AMENDMENTS TO THE CLAIMS

Amend the claims, and/or add new claims, as follows:

1. to 18. (Cancelled)

19. (Previously presented) A method for treating immunosuppression of skin cells induced by UVB radiation, said method comprising topically applying to the skin of a person in need thereof an effective amount therefor of a cosmetic or dermatological formulation comprising:

- a) one or more flavonoids selected from the group consisting of alpha-glucosylrutin, alpha-glucosylmyricitrin, alpha-glucosylisoquercitrin, alpha-glucosylquercitrin, quercitin, rutin, chrysin, kaempferol, myricetin, rhamnetin, apigenin, luteolin, naringin, hesperidin, naringenin, hesperitin, morin, phloridzin, diosmin, fisetin, vitexin, neoheperidin dihydrochalcone, flavone, glucosylrutin and genistein;
- b) optionally one or more cinnamic acid derivatives; and
- c) optionally an antioxidant;


and preventing or treating immunosuppression of skin cells induced by UVB radiation.

20. (Previously presented) The method according to claim 19, wherein the cosmetic or dermatological formulation comprises one or more cinnamic acid derivatives.


GHITA LAZENDORFER ET AL
USSN 08/849,525

21. (Previously presented) The method according to claim 20, wherein the one or more cinnamic acid derivatives are selected from the group consisting of hydroxycinnamic acid derivatives.

22. (Previously presented) The method according to claim 21, wherein the hydroxycinnamic acid derivatives are selected from those of the formula:



or the formula:




or mixtures thereof,

wherein X, Y and R independently represent H or branched or unbranched alkyl having 1 to 18 carbon atoms.


GHITA LAZENDORFER ET AL
USSN 08/849,525

23. (Previously presented) The method according to claim 19, which comprises topically applying to the skin of a person in need thereof an effective amount therefor of a cosmetic or dermatological formulation comprising:

- a) one or more flavonoids selected from the group consisting of alpha-glucosylrutin, alpha-glucosylmyricitrin, alpha-glucosylisoquercitrin, alpha-glucosylquercitrin, quercitin, rutin, chrysin, kaempferol, myricetin, rhamnetin, apigenin, luteolin, naringin, hesperidin, naringenin, hesperitin, morin, phloridzin, diosmin, fisetin, vitexin, neohesperidin dihydrochalcone, flavone, glucosylrutin and genistein;
- b) one or more hydroxycinnamic acid derivatives of the formula:



or the formula:

GHITA LAZENDORFER ET AL
USSN 08/849,525

or mixtures thereof,

wherein X, Y and R independently represent H or branched or unbranched alkyl have 1 to 18 carbon atoms; and

c) an antioxidant. --

24. (Previously presented) The method according to claim 19, wherein a symptom of the immunosuppression is inflammation, allergy or an autoimmune-reactive symptom.

25. (Previously presented) A method for protecting the cells which participate in the immune response of the skin from the damaging effects of UVB radiation, said method comprising topically applying to the skin of a person in need thereof an effective amount therefor of a cosmetic or dermatological formulation comprising:

a) one or more flavonoids selected from the group consisting of alpha-glucosylrutin, alpha-

GHITA LAZENDORFER ET AL
USSN 08/849,525

glucosylmyricitrin, alpha-glucosylisoquercitrin, alpha-glucosylquercitrin, quercitin, rutin, chrysin, kaempferol, myricetin, rhamnetin, apigenin, luteolin, naringin, hesperidin, naringenin, hesperitin, morin, phloridzin, diosmin, fisetin, vitexin, neohesperidin dihydrochalcone, flavone, glucosylrutin and genistein;


- b) optionally one or more cinnamic acid derivatives; and
- c) optionally an antioxidant;

and protecting the cells which participate in the immune response of the skin from the damaging effects of UVB radiation.


26. (Previously presented) The method according to claim 25, wherein the cosmetic or dermatological formulation comprises one or more cinnamic acid derivatives.

27. (Previously presented) The method according to claim 26, wherein the one or more cinnamic acid derivatives are selected from the group consisting of hydroxycinnamic acid derivatives.

28. (Previously presented) The method according to claim 27, wherein the hydroxycinnamic acid derivatives are selected from those of the formula:

GHITA LAZENDORFER ET AL
USSN 08/849,525

or the formula:



or mixtures thereof,

wherein X, Y and R independently represent H or branched or unbranched alkyl have 1 to 18 carbon atoms.


29. (Previously presented) The method according to claim 25, which comprises topically applying to the skin of a person in need thereof an effective amount therefor of a cosmetic or dermatological formulation comprising:

- a) one or more flavonoids selected from the group consisting of alpha-glucosylrutin, alpha-glucosylmyricitrin, alpha-glucosylisoquercitrin, alpha-


GHITA LAZENDORFER ET AL
USSN 08/849,525

glucosylquercitrin, quercitin, rutin, chrysin, kaempferol, myricetin, rhamnetin, apigenin, luteolin, naringin, hesperidin, naringenin, hesperitin, morin, phloridzin, diosmin, fisetin, vitexin, neohesperidin dihydrochalcone, flavone, glucosylrutin and genistein;

b) one or more hydroxycinnamic acid derivatives of the formula:



or the formula:



or mixtures thereof,

wherein X, Y and R independently represent H or branched or unbranched alkyl have 1 to 18 carbon atoms; and


c) an antioxidant.

GHITA LAZENDORFER ET AL
USSN 08/849,525


30. (Previously presented) A cosmetic or dermatological formulation

comprising:

- a) one or more flavonoids selected from the group consisting of alpha-glucosylrutin, alpha-glucosylmyricitrin, alpha-glucosylisoquercitrin, alpha-glucosylquercitrin, quercitin, rutin, chrysins, kaempferol, myricetin, rhamnetin, apigenin, luteolin, naringin, hesperidin, naringenin, hesperitin, morin, phloridzin, diosmin, fisetin, vitexin, neohesperidin dihydrochalcone, flavone, glucosylrutin and genistein;
- b) one or more hydroxycinnamic acid derivatives of the formula:



or the formula:

GHITA LAZENDORFER ET AL
USSN 08/849,525

or mixtures thereof,

wherein X, Y and R independently represent H or branched or unbranched alkyl have 1 to 18 carbon atoms; and

c) optionally an antioxidant.

31. (Previously presented) The cosmetic or dermatological formulation according to claim 30, which comprises an antioxidant.

32. (Previously presented) A cosmetic or dermatological formulation

comprising:

a) one or more flavonoids selected from the group consisting of alpha-glucosylmyricin, alpha-glucosylisoquercitrin, alpha-glucosylquercitrin, myricetin, rhamnetin, apigenin, hesperitin, morin, phloridzin, diosmin, vitexin, neohesperidin dihydrochalcone, flavone, and genistein;

GHITA LAZENDORFER ET AL
USSN 08/849,525

- b) optionally one or more cinnamic acid derivatives; and
- c) optionally an antioxidant.

33. (Previously presented) The cosmetic or dermatological formulation according to claim 32, which comprises an antioxidant.

34. (Previously presented) A cosmetic or dermatological formulation comprising:

- a) one or more flavonoids selected from the group consisting of alpha-glucosylmyricitrin, alpha-glucosylisoquercitrin, alpha-glucosylquercitrin, myricetin, rhamnetin, apigenin, hesperidin, hesperitin, morin, phloridzin, diosmin, vitexin, neoheperidin dihydrochalcone, flavone, glucosylrutin and genistein;
- b) one or more cinnamic acid derivatives; and
- c) one or more antioxidants.